


Maximize performance

- Cattle consume diets high in fermentable carbohydrates.
- Increases lactic acid production in the rumen, which decreases fiber digestion.
- Active, dry yeast utilize lactic acid.



Phone: (806) 632-0020 Email: info@mbfeeds.com VISIT MBFEEDS.COM FOR MORE INFORMATION









A Source of Live, Viable Saccharomyces cerevisiae - Added to the Feed of Growing Heifers and Lactating Cows

Contains a source of live, viable naturally occurring microorganisms

- Cattle consume diets high in fermentable carbohydrates
- Increases lactic acid production in the rumen, which decreases fiber digestion.
- Active, dry yeast utilize lactic acid

GUARANTEED ANALYSIS*

2 x 10¹⁰ colony forming units of Saccharomyces cerevisiae per gram

DIRECTIONS FOR USE

Lactating: 2.5-3.0 grams per head

per day

Close-up & Fresh Cows: 3 grams

per head per day

Heifers & Dry Cows: 0.5 – 1 gram

per head per day

INGREDIENTS

Active dry yeast (Saccharomyces cerevisiae)

PACKAGING & STORAGE

Available in 10 kg vacuum-sealed Mylar foil bags with 2 bags per box (#3001)

Store in a cool, dry place. Product has a shelf life of 12 months when stored in original packaging. Keep tightly closed and minimize exposure to air and moisture.



MBFEEDS.COM