



# FRESHCAL

**Do you have a Hypocalcemia problem?**

Estimates are 25-75% of cows are subclinically Hypocalcemic

**Cows suffering from Hypocalcemia experience these negative effects:**

- Reduced milk production
- Decreased conception
- Suppressed immune function
- Increased risk to be culled

**Give FRESHCAL at calving and 12 - 24 hours later**

**Each dose supplies over 50 grams of calcium from calcium chloride, sulfate, & propionate**





CALCIUM



B-GLUCAN



VIT. D<sub>3</sub>

**Estimates are 25-75% of cows are subclinically Hypocalcemic**  
**Give FRESHCAL at calving and 12 - 24 hours later**

### GUARANTEED ANALYSIS\*

Calcium (min).....50 grams  
Vitamin D3.....70,000 IU

\*Per 4 boluses

### DIRECTIONS FOR USE

#### Fresh Cow & Cow Off Feed:

Give 4 boluses immediately after calving and repeat 12 to 24 hours later if necessary.

**Primiparous Cows:** Give 4 boluses immediately after calving.

### INGREDIENTS

Calcium Chloride, Water, Calcium Sulfate, Niacin, Calcium Propionate (preservative), Yeast Extract (*Saccharomyces cerevisiae*), & Vitamin D<sub>3</sub> Supplement.

### PACKAGING & STORAGE

Individually wrapped 4 bolus dose. Available in:

- 10 count box (#5018)
- 50 count bucket (#5004)
- 90 count box (#5003)

Store bolus in original packaging at 36° - 90°F. Open sealed mylar pouch immediately before use as moisture will compromise the bolus.



**MBFEEDS.COM**